

## SPECIES

## To Cite:

Rajesekar C, Jayaganesh M, Kavitha J, Rajendran R, Kottaimuthu R.  
New record of *Brachystelma rapinatianum* Britto & Bruyns  
(Apocynaceae: Ceropegieae) from Eastern Ghats, India. *Species* 2023;  
24: e38s1530  
doi: <https://doi.org/10.54905/diss/v24i73/e38s1530>

## Author Affiliation:

<sup>1</sup>Department of Botany, Alagappa University, Karaikudi-630003,  
Tamil Nadu, India

<sup>2</sup>Presently at: Botanical Survey of India, Eastern Regional Centre,  
Shillong-793 003, Meghalaya, India

## Corresponding author

Department of Botany, Alagappa University, Karaikudi-630003,  
Tamil Nadu, India and Presently at: Botanical Survey of India,  
Eastern Regional Centre, Shillong-793 003, Meghalaya,  
India  
Email: [rkmlegumes@gmail.com](mailto:rkmlegumes@gmail.com)

## Peer-Review History

Received: 06 February 2023

Reviewed & Revised: 08/February/2023 to 20/April/2023

Accepted: 23 April 2023

Published: 27 April 2023

## Peer-Review Model

External peer-review was done through double-blind method.

## Species

pISSN 2319-5746; eISSN 2319-5754



© The Author(s) 2023. Open Access. This article is licensed under a  
Creative Commons Attribution License 4.0 (CC BY 4.0), which permits use,  
sharing, adaptation, distribution and reproduction in any medium or  
format, as long as you give appropriate credit to the original author(s) and  
the source, provide a link to the Creative Commons license, and indicate if  
changes were made. To view a copy of this license, visit  
<http://creativecommons.org/licenses/by/4.0/>

# New record of *Brachystelma rapinatianum* Britto & Bruyns (Apocynaceae: Ceropegieae) from Eastern Ghats, India

Rajesekar C<sup>1</sup>, Jayaganesh M<sup>1</sup>, Kavitha J<sup>1</sup>, Rajendran R<sup>1</sup>,  
Kottaimuthu R<sup>1,2\*</sup>

## ABSTRACT

In this communication *Brachystelma rapinatianum* Britto & Bruyns, one of the poorly known endemic geophytes is reported for the first time for the flora of Eastern Ghats from Narthamalai. Brief description, photo-plate and other relevant details are furnished.

**Keywords:** *Brachystelma rapinatianum*; Eastern Ghats; endemic geophytes

## 1. INTRODUCTION


India is considered as one of the diversity centres of *Brachystelma* and known to have 41 taxa (Paramesh et al., 2021). After the publication of *Brachystelma* of Eastern Ghats (Pullaiah et al., 2011), the continual reports of new taxa and new records (Prasad and Rao, 2013; Rasingam et al., 2013; Kullayiswamy et al., 2013, 2016; Kaliamoorthy et al., 2015; Britto and Bruyns, 2016; Prasad et al., 2016, 2017, 2018; Sadasivaiah et al., 2016; Prasad and Prasanna, 2016; Prasad and Venu, 2018, 2020; Rasingam and Swamy, 2020; Reddy et al., 2018; Nagendra et al., 2021; Paramesh et al., 2021) revealed that the data on the diversity and distribution of the genus *Brachystelma* is still incomplete in Eastern Ghats.

Floristic surveys carried out in the Narthamalai, Pudukkottai District, led to the collection of an interesting ephemeral *Brachystelma rapinatianum* Britto & Bruyns. It is hitherto known only from the plains near Tiruchirappalli (Britto and Bruyns, 2016; Britto, 2019). The species has not been collected after the type so far (Pullaiah et al., 2019; Prasad and Venu, 2020). Thus, the present collection from Narthamalai is the second report and also an addition to the flora of Eastern Ghats. A brief description and a photographic plate of the species are provided below.

## 2. TAXONOMIC TREATMENT

*Brachystelma rapinatianum* Britto & Bruyns in Haseltonia 22: 50. 2016; Pullaiah et al., (2019) Monogr. *Brachystelma* and *Ceropegia* in India: 56. 2019; K.Prasad & Venu, Taxon. Rev. of the genus *Brachystelma* in India: 77. 2020. *Ceropegia rapinatiana* (Britto & Bruyns) Bruyns in S. African J. Bot. 112: 430. 2017; Britto, Fl. Central North Tamil Nadu 3: 1800. 2019.

Erect, tuberous herbs, 10–17 cm high; tubers globose to spherical,  $2\text{--}3 \times 1.5\text{--}3$  cm, pale brown. Stem un-branched, terete, greenish purple, glandular-puberulous when young, later glabrescent. Leaves sub-sessile, opposite-decussate; lamina linear–elliptic or elliptic–oblong,  $1\text{--}1.5\text{--}2.5 \times 0.1\text{--}0.2$  cm, glandular-hairy, base acute–cuneate, margins ciliolate, apex acute–acuminate, apiculate. Inflorescence 2–5 flowered cymes; peduncle slender, up to 2 cm long, thinly glandular-puberulous; pedicels 0.5–2 cm long, slender, glandular-puberulous. Bracts linear, 1–2 mm long. Sepals 5, green, lanceolate, purplish along margins, 1 mm long, minutely glandular-puberulous. Corolla campanulate, pale green to purplish; corolla-tube ca 2 mm long, constricted near mouth; corolla lobes lanceolate-deltate, 3–4 mm long, keeled at base, erect to slightly spreading, pale green, with purplish spots, margins reflexed. Corona two-seriate; outer series forming obscure cup-like structure; inner series with 5, small, linear lobules, pale yellow, covered with purple, silky hairs. Follicles lanceolate–ellipsoid, 2–4 cm long, is spreading to erect. Seeds 8–12, ovoid, 8–10 mm across, comose at apex.



**Figure 1** *Brachystelma rapinatianum* Britto & Bruyns. a) Habit (inset tuber); b & c) Leaf; d) Inflorescence; e) Flower; f) Flower front view (inset corona); g & h) Follicle; i & j) Seed

**Flowering & Fruiting**

September–November

**Distribution**

INDIA: Tamil Nadu. Endemic.

**Specimens examined**

Tamil Nadu, Pudukkottai District, 160m, 2 Oct. 2022, Narthamalai, C. Rajasekar et al. 122 (Alagappa University Herbarium).

**Notes**

The authors have located 50 individuals at three different populations at Narthamalai. However, all these populations are rapidly dwindling mainly due to urbanization and habitat degradation along with illicit grazing. Hence further intensive explorations should be carried out in similar habitats in adjacent forests. The existing populations need to be monitored periodically. Above all, immediate efforts should be made to ensure its survival by protecting its population in the known localities and adopt ideal in-vitro propagation methods for its mass production and reintroduction at similar sites to save this endemic tuberous ephemeral from extinction.

**Acknowledgements**

Authors thank the DST-PURSE (SR/PURSE Phase 2/38 (G)) and MHRD-RUSA 2.0 (F.24/51/2014-U), Policy (TNMulti-Gen), Department of Education, Government of India for the financial assistance. We also thank the authorities of Tamil Nadu Forest Department for granting permission to conduct field studies in the reserved forests of Pudukkottai District.

**Ethical approval**

The ethical guidelines for plants & plant materials are followed in the study for sample collection & identification.

**Informed consent**

Not applicable.

**Conflicts of interests**

The authors declare that there are no conflicts of interests.

**Funding**

The study has received funding from DST-PURSE (SR/PURSE Phase 2/38 (G)) and MHRD-RUSA 2.0 (F.24/51/2014-U), Policy (TNMulti-Gen), Department of Education, Government of India.

**Data and materials availability**

All data associated with this study are present in the paper.

**REFERENCES AND NOTES**

1. Britto SJ, Bruyns PV. Three new species of *Brachystelma* from Tamil Nadu, India. *Haseltonia* 2016; 22:248–254.
2. Britto SJ. The Flora of Central and North Tamil Nadu. The Rapinat Herbarium, St. Joseph's College (Autonomous), Tiruchirappalli 2019; 3:1983.
3. Kalliamoorthy S, Saravanan TS, Elango BS, Murthy GVS. The Elenadu short crown *Brachystelma elenaduense* Sathyan. (Asclepiadaceae)-A new distribution record for Tamil Nadu, India. *J Threat Taxa* 2015; 7:7587–7588. doi: 10.11609/JoTT.o3 880.7587-8
4. Kullayiswamy R, Rani SS, Pullaiah T. *Brachystelma ciliatum* (Apocynaceae): A new record for Eastern Ghats of Andhra Pradesh, India. *Nelumbo* 2013; 55:191–195.
5. Kullayiswamy R, Rani SS, Pullaiah T. *Brachystelma nigidianum* (Apocynaceae: Asclepiadoideae), a new species from India. *Kew Bull* 2016; 71:1–7. doi: 10.1007/s12225-016-9668-z
6. Nagendra C, Nagaraju V, Reddy AM, Prasad K. A new variety of *Brachystelma maculatum* (Apocynaceae) from

- Andhra Pradesh, India. Rheedia 2021; 31:31–34. doi: 10.22244/rheedia.2021.31.01.05
7. Paramesh L, Prasad K, Sadasivaiah B, Reddy AVB. *Brachystelma ananthagiriense* (Apocynaceae), a new species from Ananthagiri hills, Telangana, India. Nord J Bot 2021; 39: 1–6. doi: 10.1111/njb.03003
  8. Prasad K, Chorghe A, Surveswaran S, Venu P. *Brachystelma mahendragiriense* (Apocynaceae), a new species from Odisha, India. Rheedia 2017; 27:135–140. doi: 10.22244/rheedia.2017.27.2.26
  9. Prasad K, Prasanna PV, Meve U, Sankara-Rao M, Thulasaih T. *Brachystelma annamacharyae* sp. nov. (Apocynaceae) from the Seshachalam hills of Andhra Pradesh (India). Nord J Bot 2016; 34:360–363. doi: 10.1111/njb.00887
  10. Prasad K, Prasanna PV. *Brachystelma seshachalamense* (Apocynaceae), a new species from Andhra Pradesh, India. Bangladesh J Plant Taxon 2016; 23:53–57.
  11. Prasad K, Rao BRP. *Brachystelma nallamalayana* sp. Nov. (Apocynaceae: Asclepiadoideae: Ceropegieae) from India. J Threat Taxa 2013; 5:4904–4906. doi: 10.11609/JoTT.o3533.4904-6
  12. Prasad K, Swamy AN, Meve U. *Brachystelma ananthapuramense* (Apocynaceae: Asclepiadoideae), a new species from Andhra Pradesh, India. Kew Bull 2018; 73:1–6. doi: 10.1007/s12225-018-9740-y
  13. Prasad K, Venu P. A taxonomic revision of the genus *Brachystelma* R.Br. in India. Bishen Singh Mahendra Pal Singh, Dehra Dun 2020.
  14. Prasad K, Venu P. Rediscovery and note on *Brachystelma maculatum* Hook. f. (Apocynaceae). Rheedia 2018; 28:73–77. doi: 10.22244/rheedia.2018.28.2.14
  15. Pullaiah T, Karuppusamy S, Ramamurthy KS. Monograph on *Brachystelma* and *Ceropegia* in India, CRC Press, Taylor & Francis Group, Boca Raton 2019.
  16. Pullaiah T, Rani S, Karuppusamy S. Flora of Eastern Ghats. *Stylidaceae to Plantaginaceae*, Regency Publication, New Delhi 2011; 4.
  17. Rasingam L, Chorghe A, Meve U, Rao MS, Prasanna PV. *Brachystelma penchalakonense* (Apocynaceae: Asclepiadoideae) – A new species from Andhra Pradesh, India. Kew Bull 2013; 68(4):663–667. doi: 10.1007/s12225-013-9477-6
  18. Rasingam L, Swamy J. *Brachystelma telanganense* (Apocynaceae: Asclepiadoideae- Ceropegieae) – a new species from Telangana, India. Rheedia 2020; 30:379–382. doi: 10.22244/rheedia.2020.30.02.02
  19. Reddy AM, Babu MVS, Prasad K. *Brachystelma vemanai* (Apocynaceae–Asclepiadoideae), a new species with completely reflexed corolla lobes and solitary flowers from Eastern Ghats, Andhra Pradesh, India. Nord J Bot 2018; 36:1–4. doi: 10.1111/njb.02067
  20. Sadasivaiah B, Kalpana B, Narasimha A, Goud MS, Rao NB. *Brachystelma nallamalayana* K. Prasad and B. R. P. Rao: A new distributional record for Telangana, India. Asian J Plant Sci Res 2016; 6:27–29.